



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Syrian immigrants inspected, 6; Mexican laborers imported in bond inspected, 107; inspection certificate of death of corpse imported in hermetically sealed box, 1; disinfection of soiled linen imported for laundry, 419 pieces; vaccination of children of immigrants, 11.

Statistical reports of States and cities of the United States—yearly and monthly.

CALIFORNIA—*Oakland*.—Month of February, 1904. Estimated population, 83,000. Total number of deaths, 77, including enteric fever 2, and 8 from tuberculosis.

CONNECTICUT.—Reports to the State board of health for the month of February, 1904, from 167 towns having an aggregate estimated population of 946,830, show a total of 1,384 deaths, including diphtheria 19, enteric fever 8, measles 9, scarlet fever 6, whooping cough 13, and 116 from phthisis pulmonalis.

FLORIDA—*Tampa*.—Month of February, 1904. Estimated population, 23,000. Total number of deaths, 37, including enteric fever 1, and 5 from tuberculosis.

ILLINOIS—*Bellefonte*.—Two weeks ended March 12, 1904. Estimated population, 19,000. Total number of deaths, 19, including enteric fever 1, and 3 from phthisis pulmonalis.

Rockford.—Month of February, 1904. Census population, 31,051. Total number of deaths, 46, including scarlet fever 1, and 4 from tuberculosis.

MICHIGAN.—Reports to the State board of health, Lansing, for the week ended March 12, 1904, from 77 observers indicate that gonorrhoea, phthisis pulmonalis, pneumonia, scarlet fever, smallpox, inflammation of bowels, remittent fever, and puerperal fever were more prevalent, and intermittent fever and meningitis were less prevalent than in the preceding week.

Meningitis was reported at 2, whooping cough at 12, diphtheria at 39, enteric fever at 44, pneumonia at 71, measles at 77, smallpox at 78, scarlet fever at 85, and phthisis pulmonalis at 214 places.

MINNESOTA—*Duluth*.—Month of February, 1904. Estimated population, 70,000. Total number of deaths, 60, including diphtheria 1, enteric fever 1, and 6 from tuberculosis.

NEW HAMPSHIRE—*Concord*.—Month of February, 1904. Estimated population, 20,000. Total number of deaths, 37, including 1 from tuberculosis.

Manchester.—Month of January, 1904. Estimated population, 60,000. Total number of deaths, 104, including diphtheria 1, enteric fever 3, whooping cough 1, and 6 from tuberculosis.

Month of February, 1904. Total number of deaths, 96, including diphtheria 1, scarlet fever 1, and 12 from tuberculosis.

NEW YORK—*Auburn*.—Month of February, 1904. Estimated population, 38,500. Total number of deaths, 36, including 6 from tuberculosis.

Buffalo.—Month of February, 1904. Estimated population, 380,000. Total number of deaths, 457, including diphtheria 9, enteric fever 5, measles 1, scarlet fever 2, whooping cough 1, and 45 from phthisis pulmonalis.

OHIO—*Cleveland*.—Month of February, 1904. Estimated population, 420,000. Total number of deaths, 682, including diphtheria 15, enteric fever 45, measles 1, and 67 from tuberculosis.

Toledo.—Month of February, 1904. Estimated population, 157,200. Total number of deaths, 134, including diphtheria 2, enteric fever 2, measles 2, and 17 from tuberculosis.

OREGON—*Portland*.—Month of February, 1904. Estimated population, 125,000. Total number of deaths, 100, including diphtheria 2, enteric fever 2, scarlet fever 4, smallpox 1, and 11 from phthisis pulmonalis.

PENNSYLVANIA—*Scranton*.—Month of February, 1904. Estimated population, 115,000. Total number of deaths, 171, including diphtheria 2, whooping cough 1, and 13 from phthisis pulmonalis.

TENNESSEE—*Knoxville*.—Month of February, 1904. Estimated population, 40,000—white, 30,000; colored, 10,000. Total number of deaths, 66—white, 44; colored, 22, including measles 2, whooping cough 2, and 15 from tuberculosis.

Nashville.—Month of February, 1904. Estimated population, 82,947; white 52,061, colored 30,886. Total number of deaths, 189; white 116, colored 73; including enteric fever 2, scarlet fever 1, and 30 from tuberculosis.

TEXAS—*San Antonio*.—Month of January, 1904. Estimated population, 65,000. Total number of deaths, 120, including diphtheria 1, and 32 from tuberculosis, of which 20 were of nonresidents.

Month of February, 1904. Total number of deaths, 156, including diphtheria 5, enteric fever 2, and 40 from tuberculosis, of which 30 were of nonresidents.

UTAH—*Salt Lake City*.—Month of February, 1904. Census population, 53,531. Total number of deaths, 111, including diphtheria 3, enteric fever 1, whooping cough 2, and 9 from tuberculosis.

VIRGINIA—*Richmond*.—Month of February, 1904. Estimated population, 100,000; white 62,250, colored 37,750. Total number of deaths, 172; white 75, colored 97; including diphtheria 2, enteric fever 1, and 24 from tuberculosis.

WASHINGTON—*Tacoma*.—Month of February, 1904. Estimated population, 65,000. Total number of deaths, 36, including 4 from phthisis pulmonalis.